Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 5 December 2012 (MN LaTeX style file v2.2)

arXiv:1212.0667v1 [astro-ph.CO] 4 Dec 2012

Dependence of Fanaroff-Riley dichotomy of radio galaxies on luminosity and redshift

Ashok K. Singal & Kamlesh Rajpurohit
Astronomy & Astrophysics Division, Physical Research Laboratory, Navrangpura, Ahmedabad - 380 009, India

E-mail: asingal@prl.res.in

Accepted . Received ; in original form

ABSTRACT

We investigate the dependence of the Fanaroff-Riley (FR) I/II dichotomy of radio galaxies on their luminosities and redshifts. Because of a very strong redshift-luminosity correlation (Malmquist bias) in a flux-limited sample, any redshift-dependent effect could appear as a luminosity related effect and vice versa. A question could then arise - do all the morphological differences seen in the two classes (FR I and II types) of sources, usually attributed to the differences in their luminosities, could these all as well be a result of mainly a cosmological evolutionary effect (e.g., due to the changing ambient density) with cosmic epoch? Even a sharp break in luminosity, seen among the two classes, could after all reflect a rather critical ambient density value. A doubt on these lines does not seem to have been raised in past and things have never been examined keeping this particular aspect in mind. We want to ascertain the customary prevalent view in the literature that the systematic differences in the two broad morphology types of FR I and II radio galaxies are indeed due to the differences in their luminosities, and not a manifestation of an evolutionary effect of the cosmic epoch. Here we investigate the dependence of FR I and II dichotomy of radio galaxies on luminosity and redshift by using the 3CR sample, where the FR I and II dichotomy was first seen, supplemented by data from two additional samples (MRC and B3-VLA), which go about a factor of 5 or more deeper in flux-density than the original 3CR sample. This lets us compare sources with similar luminosities but at different redshifts as well as examine sources at similar redshifts but with different luminosities, thereby allowing us a successful separation of the otherwise two intricately entangled effects.

Key words: galaxies: active - galaxies: nuclei - galaxies: evolution - galaxies: fundamental parameters - radio continuum: galaxies

1 INTRODUCTION

One of the robust correlations in observational astronomy is between the morphology type of radio galaxies and their radio luminosity. First pointed out by Fanaroff & Riley (1974) that there is a very sharp dependence of the morphology type of radio galaxies on their luminosity so that almost all radio galaxies below a luminosity $P_{178} = 2 \times 10^{25}$ W Hz^{-1} sr^{-1} (for Hubble constant $H_0 = 50\,\mathrm{km\ s^{-1}\,Mpc^{-1}}$), are edge-darkened (called type I) in their brightness distribution, while all radio galaxies above this luminosity limit are more or less edge-brightened (called type II). This correlation has withstood the test of time (Miley 1980; Antonucci 1993, 2012; Urry & Padovani 1995; Kembhavi & Narlikar 1999). However, because of a very strong redshift-luminosity correlation (Malmquist bias) in a flux-limited sample, like in the 3CR sample used by Fanaroff & Riley (1974), any effect related with redshift could appear as a luminosity-dependent effect and vice versa. One can then beg a question - could all differences seen in the FR I and II types of sources rather be a result of a cosmological evolutionary effect with the cosmic epoch, instead of, as almost universally believed, an effect of transition across a certain critical luminosity value?

Following the archetypal paper by Fanaroff & Riley (1974), where they first pointed out the presence of two distinct morphology types of radio galaxies in the strong source 3CR sample, ascribing the distinct morphology of each galaxy to its radio luminosity, it has ever since been thought to be only a luminosity-dependent effect (see e.g., Saripalli 2012 and the references therein). There have been no attempts to investigate the alternative possibility that it could as well be a redshift-dependent effect, demonstrat-

© 0000 RAS



Figure 1. A scatter plot of both types of FR morphologies ('Δ' for edge-darkened FR I type and '+' for edge brightened FR II type) for the 3CR source sample of Fanaroff & Riley (1974) in the redshift-luminosity plane. To avoid undue compression of the scale, we have restricted the plot to a redshift limit of 0.15. There are no FR I type radio galaxies which lie beyond a redshift 0f 0.032 or have a luminosity $P_{178} > 2 \times 10^{25}$ W Hz^{-1} sr^{-1} (for $H_0 = 50\,\mathrm{km\ s^{-1}\ Mpc^{-1}}$) in the sample of Fanaroff & Riley.

ing the strong evolution of source morphology with cosmic epoch. For instance, suppose one were to correlate the morphology type of radio galaxies with redshift. Figure 1 shows a scatter plot with redshift and luminosity of both types of morphologies classified by Fanaroff & Riley (1974) in their sample. The demarcation is as good with redshift as it is with luminosity. It might be a moot point to guess what would have been the verdict, had Fanaroff & Riley in their seminal paper tried a correlation against only redshift, instead of luminosity. It is quite likely the effect then would have been interpreted as due to a very strong evolution with cosmic epoch, and the subsequent theoretical interpretations in that case perhaps very different.

The two scenarios have very different physical interpretations, and either could be of equal importance. For instance in the conventional interpretation, with intrinsic luminosities being the root cause of their different morphology types, there is a huge amount of literature about the relation between the luminosity break and the different morphology types (Saripalli 2012 and the references therein). In fact all the models and discussions in the literature currently available are almost exclusively only within that framework. On the other hand a definite correlation with redshift alone, would imply that it is the cosmic evolution of the properties of sources (perhaps because of the ambient density falling below a certain critical value due to the Hubble expansion) that might give rise to these two different type of morphologies, with FR I type being the current favourite and the FR II type "a thing of past". All difference seen in FR I and FR II type sources, which are usually attributed to the differences in their luminosities, could as well be then related to the changing ambient densities with cosmic epoch. Also the very sharp division in luminosity could possibly be due to a critical ambient density value, which might divide the sources into two distinct morphology types. It might be the luminosity dependence or it might be the dependence on redshift that gives rise to these morphological differences, but this question could not be decisively settled based on any amount of arguments, sans actual observational data. At least this particular aspect has not yet been investigated in the literature.

For one thing, to investigate this further one needs data from different samples to disentangle the luminosity and redshift correlation in order to study the dependences of FR I and FR II break separately on them. That is to make these investigations one needs sufficient data comprising sources at different flux-density levels, so that one could examine sources with similar luminosities at different redshifts as well as compare them at similar redshifts for different luminosities, thereby separating the two effects. We investigate this dependence of FR dichotomy of radio galaxies on luminosity and redshift by taking data from three such different samples. We may add that there are reports of FR I types seen at redshifts larger than $z > 0.5$ (Saripalli et al. 2012), but a systematic investigation of this question is still needed using samples which are complete in the sense that all FR I's above the sensitivity limit of the sample are included.

2 THE SAMPLES AND THE DATA

Our first sample is the 3CR (Laing, Riley & Longair 1983), which is a complete strong source sample, with all necessary optical and radio information with good resolution maps so that one can in most cases unambiguously decide the FR I/II type of morphologies.

The second sample we have chosen is essentially complete MRC sample (Kapahi et al. 1998) with $S_{408} \geq 0.95$ Jy, which is about a factor ~ 5 deeper than the 3CR sample and has the required radio and optical information. Their total sample comprises 550 sources, with 111 of them being quasars and the remaining are radio galaxies. Optical identifications for the latter are complete up to a red magnitude of ~ 24 or a K magnitude of ~ 19 and we do not think among the remaining there would be any missing FR I types, which in any case are not expected to be seen in this sample at redshifts larger than ~ 1.

For our third sample we have chosen a homogeneous sub-sample, known as B3-VLA, from B3 sources with $S_{408} >$ 0.1 Jy, for which the relevant radio and optical information is given by Vigotti et al. (1989). Again the B3-VLA sample is expected to be complete at least as far as FR I type sources are concerned. Their VLA maps at 1415 MHz with 15 arcsec resolution are in general good enough for the morphologies to be classified into FR I/II types.

3 RESULTS AND DISCUSSION

To quantitatively distinguish between FR I and II, following Fanaroff & Riley (1974), we classify a radio galaxy as FR I if the separation between the points of peak intensity in the two lobes is smaller than half the largest size of the source. Similarly FR II is the one in which the separation between the points of peak intensity in the two lobes is greater than half the largest size of the source. This is equivalent to having the "hot spots" nearer to (FR I) or further away from (FR II) the central optical galaxy than the regions of diffuse



Figure 2. Histograms showing distributions of FR I/II morphology types radio galaxies with redshift for the (a) 3CR, (b) MRC and (c) B3-VLA samples. It should be noted that all radio galaxies seen beyond the redshift limit of the plots (i.e. $z > 0.25$) are only of FR II type in all of the above three samples.

radio emission. In all the three samples we have examined the radio maps and using the above criteria, we have classified each source into either of the two types (FR I and II). There were a small number of sources that had a doubtful classification, we have dropped them from our analysis and that should not be too detrimental to our conclusions.

Figure 2 shows histograms of the distributions of radio galaxies with FR I type morphology against redshift for all our three samples. We have restricted our plots to $z = 0.25$ only, as no FR I type radio galaxy is seen beyond this redshift in any of the three samples. We have determined median value z_{med} for the redshift distribution of FR I sources for each sample. The details of the method for determining the median values and for the estimation of their rms errors are described in Singal (1988).

Figure 3 shows similar histograms of the distribution of radio galaxies with FR I type morphology against luminosity (at 178 MHz) for our three samples. Luminosity is calculated from their flux density S_{178} and the spectral index α ($S \propto \nu^{-\alpha}$) as

$$P_{178} = S_{178}\mathcal{D}^2(1+z)^{1+\alpha}, \qquad (1)$$

where $\mathcal{D}$ is the comoving cosmological distance calculated from the cosmological redshift z of the source. In general it is not possible to express $\mathcal{D}$ in terms of z in a close-form analytical expression and one may have to evaluate it numerically. For example, in the flat universe models ($\Omega_m + \Omega_\Lambda = 1, \Omega_\Lambda \neq 0$), $\mathcal{D}$ is given by (see e.g., Weinberg 2008),

$$\mathcal{D} = \frac{c}{H_0}\int_1^{1+z} \frac{\mathrm{d}z}{(\Omega_\Lambda + \Omega_m z^3)^{1/2}}\ . \qquad (2)$$

For a given Ω_Λ, $\mathcal{D}$ can be evaluated from Eq. (2) by a numerical integration. Here Ω_m is the matter energy density (including that of the dark matter) and Ω_Λ is the vacuum energy (dark energy!) density, both defined in terms of the critical energy density $\Omega_c = 3H_0^2c^2/(8\pi G)$, where G is the gravitational constant and c is the velocity of light in vacuum. We have used $H_0 = 71\,\mathrm{km\ s^{-1}\,Mpc^{-1}}$, $\Omega_m = 0.27$ and $\Omega_\Lambda = 0.73$ (Spergel et al. 2003). Since all the FR I's we are interested in are at low redshifts ($z < 0.25$), the specific cosmological parameters do not make much difference for the luminosity evaluations except for the scaling factor due to Hubble constant, which in our case makes luminosity estimates lower by a factor of $(71/50)^2 \approx 2$ than the calculations in Fanaroff & Riley (1974). Thus the FR I/II luminosity break of $P_{178} = 2 \times 10^{25}$ W Hz^{-1} sr^{-1} (for $H_0 = 50\,\mathrm{km\ s^{-1}\,Mpc^{-1}}$), arrived at by Fanaroff & Riley



Figure 3. Histograms showing distributions of FR I/II morphology types radio galaxies with radio luminosity (P_{178} in units of W Hz^{-1} sr^{-1}) for the (a) 3CR, (b) MRC and (c) B3-VLA samples. It should be noted that all radio galaxies seen beyond the luminosity limit of the plots (i.e. $P_{178} > 10^{27}$ W Hz^{-1} sr^{-1}) are only of FR II type in all of the above three samples.

Table 1. Percentile values of redshift and luminosity distributions for the FR I sources in the three samples.

Sample	$S_{178,\mathrm{med}}$ Jy	Number FR I	z_{med}	z_{lq}	z_{uq}	$\log(P_{178,\mathrm{med}})$	$\log(P_{178,\mathrm{lq}})$	$\log(P_{178,\mathrm{uq}})$
3CR	19.4	23	0.03 ± 0.005	0.02	0.08	24.6 ± 0.2	24.2	25.5
MRC	2.7	27	0.12 ± 0.015	0.06	0.17	24.9 ± 0.1	24.6	25.2
B3-VLA	1.7	26	0.13 ± 0.015	0.06	0.16	24.7 ± 0.2	24.2	25.0

(1974) in their seminal paper, corresponds to $P_{178} = 10^{25}$ W Hz^{-1} sr^{-1} in our case. Again we have restricted our plots to $P_{178} = 10^{27}$ W Hz^{-1} sr^{-1} only, as no FR I type radio galaxy is seen beyond this luminosity value in any of the three samples.

From Figure 3, it is also clear that the FR I/II break is not as sharp as stated in Fanaroff & Riley (1974), as some overlap of both type of morphologies is seen in luminosity. However there is no denying that in all samples there are only a few, if any, FR I types with luminosities $P_{178} > 10^{26}$ W Hz^{-1} sr^{-1}. In Table 1 we have listed the median values both for the redshift and the luminosity in each of the three samples. Also listed are the median flux-density of FR I's in each sample and the number of FR I sources in the sample. The rms error in z_{med} in each case is determined from the frequency distribution (histogram) of z-distribution. The rms error is given by $\sqrt{n}/(2f_p)$ in units of the class interval of z (Kendall 1945; Yule & Kendall 1950), where n is the total number of sources in the sample and f_p is the ordinate value of the smoothened frequency distribution at the median value in Figure 2. The median value of the distribution does seem to shift substantially with redshift for samples which differ in flux-density by a factor of ~ 5 to 10. P_{med} and the rms error in P_{med} in each case is determined from the frequency distribution (histogram) of P-distribution in Figure 3, in the same way as for z_{med} as described above. Within errors there is hardly any difference in three samples in the P_{med} value, which thus seems to be independent of the flux-density level of the sample.

While the median value of redshift (z_{med}) differs for the weaker samples as much as a factor of about four, at about a 5σ level, as compared to that in the stronger 3CR sample, the difference in the median value of luminosity (P_{med}) is only marginal, being only a factor of about $10^{0.3} \sim 2$ and that too within only about 1σ. From z_{med} and $\log(P_{\mathrm{med}})$ values, it is clear that the FR I type of morphology of radio galaxies is indeed due to their luminosity below a critical value as indeed envisaged first time by Fanaroff & Riley (1974) and that it is not directly related to the redshift and hence not due to a cosmic evolution effect.

To ascertain it further we have examined the normalized cumulative distributions of FR I morphology type radio galaxies with redshift as well as luminosity. Figure 4 shows the normalized cumulative distributions of FR I morphology type radio galaxies with redshift for the three samples. From this figure we can compare the relative numbers of FR I types upto any given redshift for different samples, and thereby for different radio luminosities. If the morphology type depended only on redshift and not on luminosity, then all the three normalized cumulative distributions in Figure 4 should be more or less coincident, which definitely is not the case. In Table 1 we have listed, along with the median value (z_{med}), also the lower quartile (z_{lq}) and upper quartile (z_{uq}) of the cumulative distribution of the redshifts of the sources in the three samples. We see that the space distribution of FR I morphology type radio galaxies does change with the flux-density levels of the sample. There is a large difference in the redshift distribution of the 3CR sample from those of the MCR or B3-VLA samples, the latter two (MRC and B3-VLA samples) however, have statistically almost indistinguishable redshift distribution of the FR I radio galaxies, not surprising as they are at about similar flux-density levels, while the 3CR sample is about 5 to 10 times stronger in flux-density than the other two samples. It does seem that the redshift distribution of the FR I radio galaxies depends heavily on the flux-density level of the sample.



Figure 4. Normalized cumulative distributions of FR I morphology type radio galaxies with redshift for the (a) 3CR sample (continuous curve), (b) MRC sample (broken curves) and (c) B3-VLA sample (dashed curve). N1, N2 and N3 give the number of FR I radio galaxies in the 3CR, MRC and B3-VLA samples, respectively.

Figure 5 shows the normalized cumulative distribution of radio luminosity for FR I morphology type radio galaxies for our three samples. The three distributions overlap and there seems to be no gross difference in the luminosity distribution of FR I sources in the three samples. In Table 1 we have also listed the lower quartile (P_{lq}) and upper quartile (P_{uq}) of the cumulative distribution of the radio luminosity of the source in the three samples, which again are of very similar values for all three samples. From these we find that half of FR I types lie in the narrow range of luminosities $P_{178} \sim 10^{24.7\pm0.5}$ W Hz^{-1} sr^{-1}, that is within a factor of 3 around 5×10^{24} W Hz^{-1} sr^{-1}. Also all galaxies with radio luminosity above $P_{178} \sim 10^{26.5}$ W Hz^{-1} sr^{-1} are only of type II with an edge brightened morphology. We have thus reaffirmed that the FR I/II dichotomy is due to a change in



Figure 5. Normalized cumulative distributions of FR I morphology type radio galaxies with radio luminosity (P_{178} in units of W Hz^{-1} sr^{-1}) for the (a) 3CR sample (continuous curve), (b) MRC sample (broken curves) and (c) B3-VLA sample (dashed curve). N1, N2 and N3 give the number of FR I radio galaxies in the 3CR, MRC and B3-VLA samples, respectively.

luminosity below and above a certain critical level, as first proposed by Fanaroff & Riley (1974) and followed in the literature ever since, and that it is not due to any cosmic epoch dependent evolution that gives rise to different FR I and II type morphologies.

4 CONCLUSIONS

We have investigated if there is a direct dependence of the FR I and II morphology types of the radio galaxies on redshift. For this we compared their distributions in three different samples with different flux-density limits, which allowed us to separately study the effects of redshift and/or radio luminosity on the occurrences of the two morphology types. It was shown that the morphology type is not directly related to redshift and thereby not a cosmic epoch dependent effect. The break between the two types of morphologies seems to depend only upon the radio luminosity. Half of the FR I type radio galaxies lie in the narrow range of luminosities $P_{178} \sim 10^{24.7\pm0.5}$ W Hz^{-1} sr^{-1}, with none exceeding the value $P_{178} \sim 10^{26.5}$ W Hz^{-1} sr^{-1}, above which all were found to be exclusively type II with an edge brightened morphology.

REFERENCES

Antonucci, R., 1993, ARAA, 31, 473
Antonucci, R., 2012, astro-ph/1101.0837
Fanaroff B. L., Riley J. M., 1974, MNRAS, 167, 31P
Kapahi, V. K., Athreya, R. M., van Breugel, W., McCarthy, P. J., Subrahmanya, C. R., 1998, ApJS, 118, 275.
Kembhavi, A. K., Narlikar, J. V. 1999, *Quasars and Active Galactic Nuclei - An Introduction* CUP, Cambridge, 357
Kendall, M. G., 1945, *The Advanced Theory of Statistics*, vol. 1, p. 211, Charles Griffin & Co. Ltd., London
Laing R. A., Riley J. M., Longair M. S., 1983, MNRAS, 204, 151
Miley, G. K., 1980, ARAA, 18, 165
Saripalli, L., 2012, AJ, 144, 85
Saripalli, L., Subrahmanyan, R., Thorat, K., Ekers, R. D., Hunstead, R. W., Johnston, H. M., Sadler, E. M., 2012, ApJS, 199, 27
Singal A. K., 1988, MNRAS, 233, 87
Spergel D.N. et al., 2003, ApJS, 148, 175
Urry, C. M., Padovani, P., 1995, PASP, 107, 803
Vigotti M., Grueff G., Perley R., Clark B. G., Bridle A. H., 1989, AJ, 98, 419
Weinberg,. S., 2008, *Cosmology*, p. 43, Oxford, Oxford
Yule, U. G., Kendall, M. G., 1950, *An Introduction to the Theory of Statistics*, p. 425, Charles Griffin & Co. Ltd., London